



EXPLORABLE
Think Outside The Box

Published on *Explorable.com* (<https://explorable.com>)

[Home](#) > Sense of Smell

Sense of Smell

Sarah Mae Sincero 10.2K reads

After a long day at work or school, you finally get into a coffee shop and order a cup of coffee and some cinnamon rolls. You notice the aroma of the coffee and the sweet scent of the cinnamon rolls, and feel relaxed in an instant. This is how your sense of smell works – the airborne chemicals of the food stimulate your sensory receptors that correspond to the sense of smell.

EXPLORABLE
Quiz Time!

Quiz: Psychology 101 Part 2

Quiz: Psychology 101 Part 2

Quiz: Flags in Europe

[See all quizzes ⇒](#)


Anatomy

The smell sensory receptors are classified under a bigger group of sensory receptors known as chemoreceptors. These chemoreceptors carry out a process called transduction, wherein chemical signals from the stimulus are transformed into action potentials. Action potentials are short-lasting events that lead to the recognition of the chemical signals, and thus, the stimulus.

The smell sensory receptors are found in a tissue in the nasal cavity called the olfactory epithelium. This patch of tissue is comprised of three types of cells:

- **Basal cells** - cells that undergo regular process of division in order to produce new sensory neurons as replacement for dying neurons
- **Supporting cells** - cells located in between basal cells and sensory neurons
- **Sensory neurons** - cells paired with primary cilia and contain the olfactory receptors that receive chemical signals from the stimulus

Cilia are hair-like protuberances that are necessary for processing molecules that we smell, also known as "odorant molecules". The cilia of sensory neurons are covered with mucus where the odorant molecules are dissolved. Once dissolved, the molecules undergo a binding process which activate the G protein in the sensory receptor. When G protein is activated, the enzyme on the cilia begins to work by catalyzing the conversion of ATP (cell energy). This leads to the sodium (Na^+) electrolytes to enter the cell. Once there's an influx of sodium, the plasma membrane's potential is continuously reduced, a process called depolarization. When the depolarization of the plasma membrane reaches its limit or threshold, an action potential is generated



ory nerve to the brain. The brain
ation, and once the brain is done evaluating
perceive the particular odor.

Photo by

Zachary Veach

Different Odors

There are thousands of odors around us, yet our sense of smell [1] only includes one kind of cell. How can we differentiate one odor from another then? Many scientists agree that each olfactory receptor is able to bind to different kinds of odorants according to the odorant chemicals' intensity. For example:

Odorant Coffee binds to the olfactory receptors on neurons #7, #11, and #345.

Odorant Cinnamon binds to the olfactory receptors on neurons #7, #23, and #543.

The brain will interpret these as two different odors, and would normally correspond to one being the coffee, and the other being the cinnamon.

Source URL: <https://explorable.com/sense-of-smell>

Links

[1] <http://users.rcn.com/jkimball.ma.ultranet/BiologyPages/O/Olfaction.html>